

IOBC/WPRS Bulletin

Vol. 23(2), 2000

Working Group „Insect Pathogens and Insect Parasitic Nematodes“, Proceedings of the 7th European Meeting “Capturing the Potential of Biological Control” at Vienna (Austria), 22– 26 March 1999. Edited by: P.H. Smits. ISBN 92-9067-119-X.

Introduction	i
List of Participants	iii
Forty years of organised international insect pathology. <i>Weiser J.</i>	3
Entomopathogenic fungi in the context of pest biocontrol today. <i>Papierok B.</i>	7
A researchers' obstacle race: report on the registration of a new bioinsecticide. <i>Strasser H.</i>	9
Commercialisation of a microbial pesticide. "Challenges and constraints" <i>Pas R. K van der, C.J. Waddington & W. Ravensberg</i>	15
Present use and future potential for biological control of pests and diseases especially in organic production of field crops in Denmark. <i>Eilenberg J., A. Enkegaard & D. Funck Jensen</i>	19
Biological control of the garden chafer <i>Phyllopertha horticola</i> (L.) <i>Strasser H.</i>	23
Capturing the potential of biological control in Italy: Where are we? <i>Rovesti L. & B. Sgarzi</i>	29
Susceptibility of the cockroach, <i>Blatta orientalis</i> L. to entomopathogenic nematodes. <i>Kotlarska-Mordzinka J, W. Goszczynski & A. Bednarek</i>	39
The susceptibility of <i>Doclostaurus maroccanus</i> (Orthoptera:Acrididae) nymphs to a Spanish isolate of <i>Steinernema</i> sp. (<i>glaseri</i> group). <i>Quesada-Moraga E. & C. Santiago-Alvarez</i>	41
Pine weevil biocontrol: Compatibility of entomopathogenic nematodes and stump degrading fungi in laboratory conditions. <i>Armendáriz I., M.J. Downes & C. T. Griffin</i>	47
Preliminary study of pathogenicity of a local strain of <i>Steinernema abbasi</i> (Steinernematidae: Nematoda) to cruciferous pests. <i>Hsiao W. -F. & R. Javedan</i>	51
The influence of storage temperature and time on infectivity and reproduction of <i>Heterorhabditis megidis</i> (strain NLH-E 87.3). <i>Boff M.I.C, G. L. Wieggers & P. H. Smits</i>	53
The susceptibility of the second stage juveniles of <i>Meloidogyne javanica</i> to the bacterial symbiont from the entomopathogenic nematode, <i>Steinernema abbasi</i> (Steinernematidae: Nematoda). <i>Samaliev H. Y, F. Andreoglou, S. Elawad, N. Hague & S. Gowen</i>	61
A possible competition mechanism in insects parasitized with rhabditid nematodes making the insect carcass decay-resistant. <i>Jarosz J.</i>	67
Entomopathogenic nematodes from Steinernematidae and Heterorhabditidae (Nematoda: Rhabditida) in Bulgaria. <i>Shishiniovva M., L. Budurova & D. Gradinarov</i>	75
Nematode community structure in poplar (<i>Populus alba</i> L.) forest in eastern Croatia.	

<i>Ivezic M, E. Raspudic & M Mlinarevic</i>	79
The potential use of entomopathogenic nematodes to control ticks. <i>Samish M., E. Alekseev, & I. Glazer</i>	83
An image analysis of <i>Spodoptera littoralis</i> feeding behavior following ingestion of the insecticidal nematode <i>Steinernema riobrave</i> in an alginate gel carrier. <i>Navon A., V. Alchanatis. I. Glazer, L. Salame & S. Levski</i>	85
PCR for identifying the entomopathogenic nematode <i>Steinernema carpocapsae</i> from overwintering codling moth larvae in Bulgaria. <i>Vega F.E., A.P. Reid, F. Herard, D. Pilarska, E. Danova, R. Tomov, L.A. Lacej, & H. K. Kaya</i>	91
The role and potential of secondary metabolites of the bacterial symbionts. <i>Webster J.M.</i>	93
Insecticidal toxins from the bacterium <i>Photorhabdus luminescens</i> : gene cloning and toxin histopathology. <i>Bowen D., M. Blackburn, T.A. Rocheleau, O. Andreev, E. Golubeva & R. H. French- Constant</i>	97
Cloning and expression of insecticidal toxin genes from <i>Xenorhabdus species</i> . <i>Morgan J.A.W., P. Jarrett, D.J. Ellis, M.A. Ousley & M. Sergeant</i>	101
Quality control of entomopathogenic nematodes. <i>Grunder J.M.</i>	103
Estimation of errors in nematode counts from suspensions. <i>Kramer L</i>	105
Insect based assays for entomopathogenic nematode infectiveness: Definitions, guidelines, problems. <i>Peters A.</i>	109
Influence of insecticides, carbosulfan and carbofuran on the mortality and pathogenicity of entomopathogenic nematodes and fungus. <i>Bednarek A., E. Popowska-Nowak, E. Pezowicz & M. Kamionek</i>	115
Interactions between Steinernematid nematodes and <i>Delia radicum</i> under field conditions. <i>Nielsen O. & H. Philipson</i>	119
Do insect aggregations influence entomopathogenic nematodes occurrence ? <i>Mracek Z, S. Becvar & P. Kindlmann</i>	123
Entomophthorean fungi infecting the green spruce aphid (<i>Elatobium abietinum</i>) in the north-western part of Europe. <i>Nielsen C., J. Eilenberg, S. Harding, E. Oddsdottir & G. Halldorsson</i>	131
Advantages and disadvantages of different sampling methods for obtaining prevalence data of fungal infections in root flies. <i>Klingen L., R. Meadow & J. Eilenberg</i>	135
A molecular method to discriminate between <i>Entomophthora</i> species using frozen or alcohol preserved cadavers. <i>Bruun Jensen A. & J. Eilenberg</i>	141
Natural occurrence of entomophthorean fungi on collembolans. <i>Dromph K., J Eilenberg & P. Esbjerg</i>	147
On mycoses of phytophagous mites (Acari). <i>Mietkiewsky R., C. Tkaczuk & S. Balazy</i>	151
The pathogenicity of entomopathogenic fungi to <i>Boophilus annulatus</i> ticks. <i>Gindin G., M. Samish, E. Alekseev & L. Glazer</i>	155
Fungi associated with the coffee berry borer, <i>Hypothenemus hampei</i> . <i>Vega F. E., G. Mercadier & M.A. Jackson</i>	159
Laboratory evaluation of <i>Beauveria bassiana</i> (Bals.) Vuill. and <i>Beauveria brongniartii</i> (Sacc.) Petch against the four eyed spruce bark beetle, <i>Polygraphus poligraphus</i> (L.) (Coleoptera, Scolytidae).	

<i>Wegensteiner R.</i>	161
Preliminary investigations on the use of <i>Beauveria bassiana</i> (Bals.) Vuill. and other control methods against the bark beetle, <i>Ips typographus</i> L. (Col., Scolytidae), in the field.	
<i>Kreutz J., G. Zimmermann, H., Marohn, O. Vaupel & G. Mosbacher</i>	167
Characterization of <i>Beauveria brongniartii</i> by the BIOLOG-system.	
<i>Abendstein D., B. Pernfuss & H. Strasser</i>	175
Persistence of released <i>Metarhizium anisopliae</i> in soil and prevalence in ground and rove beetles.	
<i>Vestergaard S. & J. Eilenberg</i>	181
Entomophthorales as significant natural enemies of cereal aphids infesting maize in Slovakia.	
<i>Stalmachová M. & L. Cagán</i>	187
<i>Cordyceps militaris</i> (Fr.) Link. in the outbreak of <i>Dendrolimus pini</i> L. in Lithuania.	
<i>Gedminas A., P. Zolubas & A. Ziogas</i>	193
Temperature as a selective factor for isolation of entomopathogenic fungi from soil by means of the insect bait method.	
<i>Tkaczuk C., R. Mietkiewski & S. Balazy</i>	197
Estimation of virulence by measuring the chitinase activity in submerged culture of <i>Beauveria brongniartii</i> and <i>Metarhizium anisopliae</i> .	
<i>Jensen D.B. & H. Strasser</i>	203
Effect of magnesium ions on the pathogenicity of entomopathogenic microorganisms applied into contaminated soil.	
<i>Ropek D. & A. Gorczyca</i>	207
Accumulation of some metal ions on biomass of insect pathogenic fungus <i>Paecilomyces fumosoroseus</i> .	
<i>Gorczyca A. & D. Ropek</i>	211
<i>Paecilomyces fumosoroseus</i> : Efficacy against the cotton pests <i>Spodoptera littoralis</i> (Lepidoptera: Noctuidae), <i>Helicoverpa armigera</i> (Lepidoptera: Noctuidae) and <i>Aphis gossypii</i> (Homoptera: Aphididae) and effect of temperature on germination and mycelial growth.	
<i>Hassani M., G. Zimmermann & S. Vidal</i>	217
Importance of coincidence for the efficiency of the entomopathogenic fungus <i>Verticillium lecanii</i> against <i>Frankliniella occidentalis</i> .	
<i>Sermann H.</i>	223
Investigations on the adhesion of spores of <i>Verticillium lecanii</i> at <i>Frankliniella occidentalis</i> (Pergande, 1895) (Thysanoptera: Thripidae).	
<i>Wolff N.B., H. Sermann & H. Bochow</i>	227
Pathogenicity of several entomogenous fungi to some of the most serious forest insect pests in Europe.	
<i>Markova, G.</i>	231
Preliminary results of experiments for the use of baculoviruses in Polish forestry.	
<i>Skrzecz L.</i>	243
Persistence and accumulation of <i>Malacosoma neustria</i> nuclear polyhedrosis virus in the ecosystem after virus application.	
<i>Jankevica L.</i>	249
Susceptibility of newly emerged adults of <i>Leptinotarsa decemlineata</i> (Say) to <i>Bacillus thuringiensis</i> ssp. <i>tenebrionis</i> .	
<i>Balasi G. & G. -A. Langenbruch</i>	255
BIOBIT XL (<i>Bacillus thuringiensis</i> Berliner subsp. <i>kurstaki</i>) in controlling ECB (<i>Ostrinia nubilalis</i> Hübner) on corn in East Croatia.	
<i>Raspudic E., M. Ivezic & M. Minarevic</i>	261

Expression of <i>cryIVA</i> and <i>cryIVD</i> genes of <i>Bacillus thuringiensis</i> in baculovirus expression system. Demir I., R. Nalcacioglu, Z. Demirbag, A. O. Kilic & A. O. Beldüz	267
New mosquitocidal <i>Bacillus thuringiensis</i> strains and non larvicidal <i>Bacillus thuringiensis</i> var. <i>israelensis</i> strains. Lage J., I. Thiéry, J. Bresciani & J. Eilenberg	275
Natural occurrence of <i>Bacillus thuringiensis</i> in Lithuanian forest ecosystems. Thomsen L., J. Eilenberg, P. Zolubas, A. Ziogas & S. Harding	279
Evaluation of the potential of a microsporidian isolate for the biological control of <i>Lymantria dispar</i> L. Linde A., J. Gollmack & D. Goertz	285
Preliminary results on the occurrence of microsporidia of gipsy moth (<i>Lymantria dispar</i> L.) from different forest habitats of Poland. Sierpinska A.	291
Study of gypsy moth microsporidia transmission by <i>Glyptapanteles liparidis</i> and <i>Cotesia melanoscela</i> (Hym.: Braconidae). Zubrik M. & J. Novotny	297
Morphometrical comparison of various microsporidian isolates from the gypsy moth, <i>Lymantria dispar</i> L. Pilarska D., A. Linde & P. Pilarski	303
Diagnosis of insect pathogens from locusts and other orthopterans. Kleespies R. G. & A.M. Huger	311