

IOBC/WPRS Bulletin

Vol. 23(8) 2000

Working Group "Integrated control of soil pests, *Melolontha* Subgroup". Proceedings of the Meeting at Sion (Switzerland), 19 – 21 October 1998. Edited by: S. Keller. ISBN 92-9067-124-6.

Introduction	i
List of participants	iii
The evolution of the situation about the <i>Melolontha melolontha</i> in Aosta valley <i>Bondaz F.</i>	3
The common cockchafer (<i>Melolontha melolontha</i> L.) in the Adige valley (Trentino): development of the population from 1987 - 1998 <i>Mattedi L. & Varner M.</i>	5
Cockchafer (<i>Melolontha melolontha</i>) - The last 20 years in the Valais <i>Schmid A.</i>	11
Possibility of the integrated control of <i>Melolontha</i> larvae <i>Bednarek M., Popowska E., Pezowicz E., Kamionek M. & Malinowski H.</i>	15
Integrated method to control <i>Melolontha</i> spp. larvae and adults in Poland <i>Stocki J.S. & Malinowski H.</i>	19
The 1997 control campaign of <i>Melolontha melolontha</i> L. at the Kaiserstuhl area (Baden- Württemberg) Field trials and practical experiences <i>Fröschle M. & Glas M.</i>	27
Potential for microbial control of scarabs and weevils in Danish forestry <i>Vestergaard S., Eilenberg J. & Harding S.</i>	35
Susceptibility of <i>Melolontha melolontha</i> to <i>Heterorhabditis bacteriophora</i> , <i>H. megidis</i> and <i>Steinernema glaseri</i> <i>Peters A.</i>	39
Investigations on the infection biology of <i>Bacillus popilliae</i> <i>Krieger L., Zhang J. & Schnetter W.</i>	47
Why is the scarab specific <i>Bacillus thuringiensis</i> ssp <i>japonensis</i> strain <i>bulbui</i> inefficient against <i>Melolontha</i> spec. <i>Wagner W., Krieger L. & Schnetter W.</i>	55
Investigations on biological control of <i>Melolontha hippocastani</i> : Results on research activities 1997 <i>Zimmermann G., Kleespies R.G. & Bathon H.</i>	61
Use of <i>Beauveria brongniartii</i> in Switzerland and its acceptance by farmers <i>Keller S.</i>	67
Insect pathogenic fungi from <i>Melolontha melolontha</i> control sites in the canton Thurgau <i>Keller S., David-Henriet A-I., & Schweizer C.</i>	73
RAPD derived markers: A rapid method for identifying <i>Beauveria brongniartii</i> <i>Piatti P., Cravanzola F., Ozino O.I. & Bondaz F.</i>	79
Field application of <i>Beauveria brongniartii</i> , established on peeled barley kernels (remarks to a field demonstration) <i>Matzke H.</i>	87
Progress report on the registration of <i>Beauveria brongniartii</i> <i>Strasser H.</i>	93
Considerations on "toxic metabolites produced by <i>Beauveria brongniartii</i> <i>Abendstein D. & Strasser H.</i>	99
Estimating chitinase activity of <i>Beauveria brongniartii</i> in submerged culture <i>Lung T., Strasser H. & Schinner F.</i>	107

Oosporein, a fungal secondary metabolite with antimicrobial properties <i>Strasser H. & Abendstein D.</i>	113
Melolonthine pests of sugarcane in Australia: An overview of research and management <i>Logan D.P. & Allsopp, P.G.</i>	119
Review of the development of <i>Metarhizium anisopliae</i> as a microbial insecticide, BioCane™, for the control of greyback canegrub <i>Dermolepida albohirtum</i> (Waterhouse) (Coleoptera: Scarabaeidae) in Queensland sugarcane <i>Logan D.P., Robertson L.N. & Milner R.J.</i>	131